

# References

## Chapter 2

- 1 Pope C, Mays N. Reaching the parts other methods cannot reach: an introduction to qualitative methods in health and health services research. *British Medical Journal (Clinical Research Edition)*. 1995;**311**:42–5.
- 2 Grimes DA, Schulz KF. Descriptive studies: what they can and cannot do. *Lancet*. 2002;**359**:145–9.
- 3 Wettermark B, Nyman K, Bergman U. Five years' experience of quality assurance and feedback with individual prescribing profiles at a primary health-care centre in Stockholm, Sweden. *Quality in Primary Care*. 2004;**12**:225–34.
- 4 Sermet C, Andrieu V, Godman B, Van Ganse E, Haycox A, Reynier JP. Ongoing pharmaceutical reforms in France: implications for key stakeholder groups. *Applied Health Economics and Health Policy*. 2010;**8**:7–24.
- 5 Garuoliene K, Godman B, Gulbinovic J, Wettermark B, Haycox A. European countries with small populations can obtain low prices for drugs: Lithuania as a case history. *Expert Review of Pharmacoeconomics & Outcomes Research*. 2011;**11**:343–9.
- 6 Lupattelli A, Spigset O, Twigg MJ, et al. Medication use in pregnancy: a cross-sectional, multinational web-based study. *British Medical Journal Open*. 2014;**4**:e004365.
- 7 Languasco A, Galante M, Marin J, Soler C, Lopez Saubidet C, Milberg M. Adherence to local guidelines for venous thromboprophylaxis: a cross-sectional study of medical inpatients in Argentina. *Thrombosis Journal* 2011;**9**:18.
- 8 Elseviers MM, Vander Stichele RR, Van Bortel L. Quality of prescribing in Belgian nursing homes: an electronic assessment of the medication chart. *International Journal for Quality in Health Care*. 2014;**26**:93–9.
- 9 Galvao TF, Silva MT, Gross R, Pereira MG. Medication use in adults living in Brasilia, Brazil: a cross-sectional, population-based study. *Pharmacoepidemiology and Drug Safety*. 2014;**23**:507–14.
- 10 Dong L, Wang D, Gao J, Yan H. Doctor's injection prescribing and its correlates in village health clinics across 10 provinces of Western China. *Journal of Public Health (Oxford, England)*. 2011;**33**:565–70.
- 11 Lusini G, Lapi F, Sara B, et al. Antibiotic prescribing in paediatric populations: a comparison between Viareggio, Italy and Funen, Denmark. *European Journal of Public Health*. 2009;**19**:434–8.
- 12 Loikas D, Wettermark B, von Euler M, Bergman U, Schenck-Gustafsson K. Differences in drug utilisation between men and women: a cross-sectional analysis of all dispensed drugs in Sweden. *British Medical Journal Open*. 2013;**3**.
- 13 Watkins C, Harvey I, Carthy P, Moore L, Robinson E, Brawn R. Attitudes and behaviour of general practitioners and their prescribing costs: a national cross sectional survey. *Quality & Safety in Health Care*. 2003;**12**:29–34.
- 14 Melander A, Folino-Gallo P, Walley T, et al. Utilisation of antihyperglycaemic drugs in ten European countries: different developments and different levels. *Diabetologia*. 2006;**49**:2024–9.
- 15 Hollingworth SA, Rush A, Hall WD, Eadie MJ. Utilization of anti-Parkinson drugs in Australia: 1995–2009. *Pharmacoepidemiology and Drug Safety*. 2011;**20**:450–6.
- 16 Davidsen JR, Sondergaard J, Hallas J, et al. Impact of socioeconomic status on the use of inhaled corticosteroids in young adult asthmatics. *Respiratory Medicine*. 2011;**105**:683–90.
- 17 Boyc KS, Yurgin N, Lage MJ. Trends in the prescription of antidiabetic medications in France: evidence from primary care physicians. *Advances in Therapy*. 2007;**24**:803–13.
- 18 Zoega H, Baldursson G, Hrafnkelsson B, Almarsdottir AB, Valdimarsdottir U, Halldorsson M. Psychotropic drug use among Icelandic children: a nationwide

- population-based study. *Journal of Child and Adolescent Psychopharmacology*. 2009;**19**:757–64.
- 19 Kotwani A, Holloway K. Trends in antibiotic use among outpatients in New Delhi, India. *BMC Infectious Diseases*. 2011;**11**:99.
- 20 Mojtabai R, Olfson M. National trends in long-term use of antidepressant medications: results from the US National Health and Nutrition Examination Survey. *The Journal of Clinical Psychiatry*. 2014;**75**:169–77.
- 21 Leri F, Ayzenberg M, Voyce SJ, Klein A, Hartz L, Smego RA Jr. Four-year trends of inappropriate proton pump inhibitor use after hospital discharge. *Southern Medical Journal*. 2013;**106**:270–3.
- 22 Vitry A, Lai YH. Advertising of antihypertensive medicines and prescription sales in Australia. *Internal Medicine Journal* 2009;**39**:728–32.
- 23 Vozoris NT, Fischer HD, Wang X, et al. Benzodiazepine use among older adults with chronic obstructive pulmonary disease: a population-based cohort study. *Drugs & Aging*. 2013;**30**:183–92.
- 24 Teuner CM, Menn P, Heier M, Holle R, John J, Wolfenstetter SB. Impact of BMI and BMI change on future drug expenditures in adults: results from the MONICA/KORA cohort study. *BMC Health Services Research*. 2013;**13**:424.
- 25 Karlsson L, Wettermark B, Tomson T. Drug treatment in patients with newly diagnosed unprovoked seizures/epilepsy. *Epilepsy Research* 2014;**108**:902–8.
- 26 Citarella A, Kieler H, Sundstrom A, et al. Family history of cardiovascular disease and influence on statin therapy persistence. *European Journal of Clinical Pharmacology*. 2014;**70**:701–7.
- 27 Sino CG, Stuffken R, Heerdink ER, Schuurmans MJ, Souverein PC, Egberts TA. The association between prescription change frequency, chronic disease score and hospital admissions: a case control study. *BMC Pharmacology & Toxicology*. 2013;**14**:39.
- 28 Burr NE, Talboys RJ, Savva S, et al. Aspirin may prevent cholangiocarcinoma: a case-control study from the United Kingdom. *Digestive Diseases and Sciences*. 2014;**59**:1567–72.
- 29 Willson MN, Greer CL, Weeks DL. Medication regimen complexity and hospital readmission for an adverse drug event. *The Annals of Pharmacotherapy*. 2014;**48**:26–32.
- 30 Breekveldt-Postma NS, Erkens JA, Aalbers R, van de Ven MJ, Lammers JW and Herings RM. Extent of uncontrolled disease and associated medical costs in severe asthma – a PHARMO study. *Current Medical Research and Opinion*. 2008;**24**:975–83.
- 31 van der Hooft CS, Schoofs MW, Ziere G, et al. Inappropriate benzodiazepine use in older adults and the risk of fracture. *British Journal of Clinical Pharmacology*. 2008;**66**:276–82.
- 32 Juurlink DN, Mamdani M, Kopp A, Laupacis A, Redelmeier DA. Drug-drug interactions among elderly patients hospitalized for drug toxicity. *Journal of the American Medical Association*. 2003;**289**:1652–8.
- 33 Grimshaw J, Campbell M, Eccles M, Steen N. Experimental and quasi-experimental designs for evaluating guideline implementation strategies. *Family Practice* 2000;**17**(Suppl. 1):S11–16.
- 34 Stephenson J, Imrie J. Why do we need randomised controlled trials to assess behavioural interventions? *British Medical Journal (Clinical Research Edition)*. 1998;**316**:611–13.
- 35 O'Connell DL, Henry D, Tomlins R. Randomised controlled trial of effect of feedback on general practitioners' prescribing in Australia. *British Medical Journal (Clinical Research Edition)*. 1999;**318**:507–11.
- 36 Lu CY, Ross-Degnan D, Stephens P, Liu B, Wagner AK. Changes in use of antidiabetic medications following price regulations in China (1999–2009). *Journal of Pharmaceutical Health Services Research*. 2013;**4**:3–11.
- 37 Sondergaard J, Foged A, Kragstrup J, et al. Intensive community pharmacy intervention had little impact on triptan consumption: a randomized controlled trial. *Scandinavian Journal of Primary Health Care*. 2006;**24**:16–21.
- 38 Martikainen JE, Saastamoinen LK, Korhonen MJ, Enlund H, Helin-Salmivaara A. Impact of restricted reimbursement on the use of statins in Finland: a register-based study. *Medical Care*. 2010;**48**:761–6.
- 39 van Doormaal JE, van den Bemt PM, Zaal RJ, et al. The influence that electronic prescribing has on medication errors and preventable adverse drug events: an interrupted time-series study. *Journal of the American Medical Informatics Association*. 2009;**16**:816–25.
- 40 "Pettersson E, Vernby Å, Mölstad S, Stålsby Lundborg C. Can a multifaceted intervention targeting both nurses and physicians change the prescribing of antibiotics to nursing home residents? A cluster randomized controlled trial. *Journal of Antimicrobial Chemotherapy*. 2011;**66**(11):2659–66." Renumber all subsequent references accordingly.

- 41 Lambert MF, Masters GA, Brent SL. Can mass media campaigns change antimicrobial prescribing? A regional evaluation study. *Journal of Antimicrobial Chemotherapy*. 2007;**59**:537–43.
- 42 Dormuth CR, Carney G, Taylor S, Bassett K, Maclure M. A randomized trial assessing the impact of a personal printed feedback portrait on statin prescribing in primary care. *Journal of Continuing Education in the Health Professions*. 2012;**32**:153–62.
- 43 Lipsey MW, Wilson DB. The efficacy of psychological, educational, and behavioral treatment. Confirmation from meta-analysis. *The American Psychologist*. 1993;**48**:1181–209.
- 44 Cochrane Effective Practice and Organisation of Care Review Group. Data collection checklist. Available from: <http://epoc.cochrane.org/sites/epoc.cochrane.org/files/uploads/datacollectionchecklist.pdf> (last accessed 18 November 2015).
- 45 Grimshaw JM, Thomas RE, MacLennan G, et al. Effectiveness and efficiency of guideline dissemination and implementation strategies. *Health Technology Assessment (Winchester, England)*. 2004;**8**:iii–iv,1–72.
- 46 Wagner AK, Soumerai SB, Zhang F, Ross-Degnan D. Segmented regression analysis of interrupted time series studies in medication use research. *Journal of Clinical Pharmacy and Therapeutics*. 2002;**27**:299–309.
- 47 Morgenstern H. Ecologic studies in epidemiology: concepts, principles, and methods. *Annual Review of Public Health*. 1995;**16**:61–81.
- 48 Bronzwaer SL, Cars O, Buchholz U, et al. A European study on the relationship between antimicrobial use and antimicrobial resistance. *Emerging Infectious Diseases*. 2002;**8**:278–82.
- 49 Carlsten A, Waern M, Ekedahl A, Ranstam J. Anti-depressant medication and suicide in Sweden. *Pharmacoepidemiol Drug Safety*. 2001;**10**(6):525–30.
- 50 Gusmao R, Quintao S, McDaid D, et al. Antidepressant utilization and suicide in Europe: An ecological multi-national study. *PLoS ONE*. 2013;**8**:e66455.
- 51 Kwong JC, Maaten S, Upshur RE, Patrick DM, Marra F. The effect of universal influenza immunization on antibiotic prescriptions: an ecological study. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America*. 2009;**49**:750–6.
- 52 Xu Y, Bentley RJ, Kavanagh AM. Gender equity and contraceptive use in China: an ecological analysis. *Women & Health*. 2011;**51**:739–58.
- 53 Vancheri F, Wettermark B, Strender LE, Backlund LG. Trends in coronary heart disease mortality and statin utilization in two European areas with different population risk levels: Stockholm and Sicily. *International Cardiovascular Forum Journal*. 2014;**3**:140–6.
- 54 Lee YK, Ha YC, Park C, Yoo JJ, Shin CS, Koo KH. Bisphosphonate use and increased incidence of subtrochanteric fracture in South Korea: results from the National Claim Registry. *Osteoporosis International*. 2013;**24**:707–11.
- 55 Sofianopoulou E, Rushton SP, Diggle PJ, Pless-Mulloli T. Association between respiratory prescribing, air pollution and deprivation, in primary health care. *Journal of Public Health (Oxford, England)*. 2013;**35**:502–9.
- 56 Martin RM, Wheeler BW, Metcalfe C, Gunnell D. What was the immediate impact on population health of the recent fall in hormone replacement therapy prescribing in England? Ecological study. *Journal of Public Health (Oxford, England)*. 2010;**32**:555–64.
- 57 Kozman D, Graziul C, Gibbons R, Alexander GC. Association between unemployment rates and prescription drug utilization in the United States, 2007–2010. *BMC Health Services Research*. 2012;**12**:435.
- 58 Rikala M, Hartikainen S, Sulkava R, Korhonen MJ. Validity of the Finnish Prescription Register for measuring psychotropic drug exposures among elderly finns: a population-based intervention study. *Drugs & Aging*. 2010;**27**:337–49.
- 59 Dahlen E, Almqvist C, Bergstrom A, Wettermark B, Kull I. Factors associated with concordance between parental-reported use and dispensed asthma drugs in adolescents: findings from the BAMSE birth cohort. *Pharmacoepidemiology and Drug Safety*. 2014;**23**:942–9.
- 60 Capella D, Pedros C, Vidal X, Laporte JR. Case-population studies in pharmacoepidemiology. *Drug Safety*. 2002;**25**:7–19.
- 61 Theophile H, Laporte JR, Moore N, Martin KL, Begaud B. The case-population study design: an analysis of its application in pharmacovigilance. *Drug Safety*. 2011;**34**:861–8.
- 62 Roujeau JC, Guillaume JC, Fabre JP, Penso D, Flechet ML, Girre JP. Toxic epidermal necrolysis (Lyell syndrome). Incidence and drug etiology in France, 1981–1985. *Archives of Dermatology*. 1990;**126**:37–42.

- 63** Roujeau JC, Kelly JP, Naldi L, et al. Medication use and the risk of Stevens–Johnson syndrome or toxic epidermal necrolysis. *New England Journal of Medicine*. 1995;**333**:1600–7.
- 64** Mockenhaupt M, Viboud C, Dunant A, et al. Stevens–Johnson syndrome and toxic epidermal necrolysis: assessment of medication risks with emphasis on recently marketed drugs. The EuroSCAR-study. *Journal of Investigative Dermatology*. 2008;**128**:35–44.
- 65** Hallas J. Evidence of depression provoked by cardiovascular medication: a prescription sequence symmetry analysis. *Epidemiology (Cambridge, MA)*. 1996;**7**:478–84.
- 66** Hersom K, Neary MP, Levaux HP, Klaskala W, Strauss JS. Isotretinoin and antidepressant pharmacotherapy: a prescription sequence symmetry analysis. *Journal of the American Academy of Dermatology*. 2003;**49**:424–32.
- 67** Garrison SR, Dormuth CR, Morrow RL, Carney GA, Khan KM. Nocturnal leg cramps and prescription use that precedes them: a sequence symmetry analysis. *Archives of Internal Medicine*. 2012;**172**:120–6.
- 68** Lai EC, Hsieh CY, Kao Yang YH, Lin SJ. Detecting potential adverse reactions of sulpiride in schizophrenic patients by prescription sequence symmetry analysis. *PLoS ONE*. 2014;**9**:e89795.
- 69** von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet*. 2007;**370**:1453–7.
- 70** Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *International Journal for Quality in Health Care*. 2007;**19**:349–57.
- 71** Langan SM, Benchimol EI, Guttman A, et al. Setting the RECORD straight: developing a guideline for the Reporting of studies Conducted using Observational Routinely collected Data. *Clinical epidemiology*. 2013;**5**:29–31.
- 72** European Network of Centres for Pharmacoepidemiology and Pharmacovigilance. *ENCePP Guide on Methodological Standards in Pharmacoepidemiology*. ENCePP, 2014.
- 73** Epstein M. Guidelines for good pharmacoepidemiology practices (GPP). *Pharmacoepidemiology and Drug Safety*. 2005;**14**:589–95.